

NONCOMMUTATIVE GEOMETRY AND DUAL COALGEBRAS

LIEVEN LE BRUYN

ABSTRACT. In arXiv:math/0606241v2 M. Kontsevich and Y. Soibelman argue that the category of noncommutative (thin) schemes is equivalent to the category of coalgebras. We propose that under this correspondence the affine scheme $\text{rep}(A)$ of a k -algebra A is the dual coalgebra A° and draw some consequences. In particular, we describe the dual coalgebra A° of A in terms of the A_∞ -structure on the Yoneda-space of all the simple finite dimensional A -representations.

CONTENTS

1. $\text{rep}(A) = A^\circ$	1
2. $\text{simp}(A) = \text{corad}(A^\circ)$	2
3. the dual coalgebra A°	4
References	5

1. $\text{rep}(A) = A^\circ$

Throughout, k will be a (commutative) field with separable closure $\bar{k}$. In [3, §I.2] Maxim Kontsevich and Yan Soibelman define a *noncommutative thin scheme* to be a covariant functor commuting with finite projective limits

$$X : \text{alg}_k^{fd} \longrightarrow \text{sets}$$

from the category alg_k^{fd} of all *finite dimensional* k -algebras (associative with unit) to the category sets of all sets. They prove [3, Thm. 2.1.1] that every noncommutative thin scheme is represented by a k -coalgebra.

Recall that a k -coalgebra is a k -vectorspace C equipped with linear structural morphisms: a comultiplication $\Delta : C \longrightarrow C \otimes C$ and a counit $\epsilon : C \longrightarrow k$ satisfying the coassociativity $(id \otimes \Delta)\Delta = (\Delta \otimes id)\Delta$ and counitary property $(id \otimes \epsilon)\Delta = (\epsilon \otimes id)\Delta = id$.

By being representable they mean that every noncommutative thin scheme X has associated to it a k -coalgebra C_X with the property that for any finite dimensional k -algebra B there is a natural one-to-one correspondence

$$X(B) = \text{alg}_k(B, C_X^*)$$

Here, for a k -coalgebra C we denote by C^* the space of linear functionals $\text{Hom}_k(C, k)$ which acquires a k -algebra structure by dualizing the structural coalgebra morphisms.

They call C_X the *coalgebra of distributions* on X and define the *noncommutative algebra of functions* on X to be the dual k -algebra $k[X] = C_X^*$.

Whereas the dual C^* of a k -coalgebra is always a k -algebra, for a k -algebra A it is not true in general that the dual vectorspace A^* is a coalgebra, because $(A \otimes A)^* \not\cong A^* \otimes A^*$. Still, one can define the subspace

$$A^\circ = \{f \in A^* = \text{Hom}_k(A, k) \mid \ker(f) \text{ contains a two-sided ideal of finite codimension}\}$$

and show that the duals of the structural morphisms on A determine a k -coalgebra structure on this dual coalgebra A° , see for example [5, Prop. 6.0.2].

With these definitions, *Kostant duality* asserts that the functors

$$\text{alg}_k \begin{array}{c} \xrightarrow{o} \\ \xleftarrow{*} \end{array} \text{coalg}_k$$

are adjoint, [5, Thm. 6.0.5]. That is, for any k -algebra A and any k -coalgebra C , there is a natural one-to-one correspondence between the homomorphisms

$$\text{alg}_k(A, C^*) = \text{coalg}_k(C, A^o)$$

Moreover, we have [5, Lemma 6.0.1] that for $f \in \text{alg}_k(A, B)$, the dual map f^* determines a k -coalgebra morphism $f^* \in \text{coalg}_k(B^o, A^o)$.

For a k -algebra A one can define the contravariant functor $\text{rep}(A)$ describing its finite dimensional representations [3, Example 2.1.9]

$$\text{rep}(A) : \text{coalg}_k^{fd} \longrightarrow \text{sets} \quad C \mapsto \text{alg}_k(A, C^*)$$

from *finite dimensional k -coalgebras* coalg_k^{fd} to *sets*, which commutes with finite direct limits. As on finite dimensional k -(co)algebras Kostant duality is an anti-equivalence of categories

$$\text{alg}_k^{fd} \begin{array}{c} \xrightarrow{*} \\ \xleftarrow{*} \end{array} \text{coalg}_k^{fd}$$

we might as well describe $\text{rep}(A)$ as the noncommutative thin scheme represented by A^o

$$\text{rep}(A) : \text{alg}_k^{fd} \longrightarrow \text{sets} \quad B = C^* \mapsto \text{alg}_k(A, B = C^*) = \text{coalg}_k(C = B^*, A^o)$$

the latter equality follows again from Kostant duality. Therefore, we propose

Definition 1. *The noncommutative affine scheme $\text{rep}(A)$ is the noncommutative (thin) scheme corresponding to the dual k -coalgebra A^o of A .*

$$2. \text{simp}(A) = \text{corad}(A^o)$$

The dual k -coalgebra A^o is usually a *huge* object and hence contains a lot of information about the k -algebra A . Let us begin by recalling how the geometry of a commutative affine k -scheme X is contained in the dual coalgebra A^o of its coordinate ring $A = \mathbb{k}[X]$.

Recall that a coalgebra D is said to be *simple* if it has no proper nontrivial subcoalgebras. In particular, a simple coalgebra D is finite dimensional over k and by duality is such that D^* is a simple k -algebra, that is, D^* is a central simple L -algebra where L is a finite separable extension of k .

Hence, in case $A = \mathbb{k}[X]$ (and $\mathbb{k}$ is separably closed) we have that all simple subcoalgebras of A^o are one-dimensional (and hence are spanned by a group-like element), because they correspond to simple representations of A .

That is, A^o is *pointed* and by [5, Prop. 8.0.7] we know that any cocommutative pointed coalgebra is the direct sum of its *pointed irreducible components* (at the algebra level, this says that a semi-local commutative algebra is the direct sum of locals). Therefore,

$$A^o = \bigoplus_{x \in X} C_x$$

where each C_x is pointed irreducible and cocommutative. As such, each C_x is a subcoalgebra of the enveloping coalgebra of the abelian Lie algebra on the tangent space $T_x(X)$. That is, we recover the points of X as well as tangent information from the dual coalgebra A^o .

But then, the dual algebra of A^o , that is the 'noncommutative' algebra of functions A^{o*} decomposes as

$$A^{o*} = \bigoplus_{x \in X} \hat{\mathcal{O}}_{x, X}$$

the direct sum of the completions of the local algebras at points. The diagonal embedding $A = \mathbb{k}[X] \hookrightarrow A^{o*}$ inevitably leads to the structure sheaf $\mathcal{O}_X$.

We will now associate a topological space associated to any k -algebra A , generalizing the space of points equipped with the Zariski topology when A is a commutative affine

k -algebra. In the next section we will describe the dual coalgebra A° when A is a noncommutative affine $\mathbb{k}$ -algebra.

The *coradical* $\text{corad}(C)$ of a k -coalgebra C is the (direct) sum of all simple subcoalgebras of C . It is also the direct sum of all simple subcomodules of C , when C is viewed as a left (or right) C -comodule.

In the example above, when $A = \mathbb{k}[X]$, we have that $\text{corad}(A^\circ) = \bigoplus_{x \in X} \mathbb{k} \, ev_x$ where the group-like element ev_x is evaluation in the point x . This motivates :

Definition 2. For a k -algebra A we define the space of points $\text{simp}(A)$ to be the set of direct summands of $\text{corad}(\text{rep}(A)) = \text{corad}(A^\circ)$. That is, $\text{simp}(A)$ is the set of simple subcoalgebras of $\text{rep}(A)$.

By Kostant duality it follows that $\text{simp}(A)$ is the set of all finite dimensional simple algebra quotients of the k -algebra A , or equivalently, the set of all isomorphism classes of finite dimensional simple A -representations, explaining the notation.

We can equip this set with a *Zariski topology* in the usual way, using the *evaluation map*

$$A^\circ \times A \xrightarrow{ev} k \quad (f, a) \mapsto f(a)$$

when restricted to the subcoalgebra $\text{corad}(A^\circ)$. Note that the evaluation map actually defines a *measuring* of A to k [5, Prop. 7.0.3], that is, $A^\circ \otimes A \xrightarrow{ev} k$ satisfies

$$ev(f \otimes aa') = \sum_{(f)} f_{(1)}(a) f_{(2)}(a') \quad \text{and} \quad ev(f \otimes 1) = \epsilon(f) 1_k$$

Definition 3. The Zariski topology of a k -algebra A is the set $\text{simp}(A)$ equipped with the topology generated by the basic closed sets

$$\mathbb{V}(a) = \{S \in \text{simp}(A) \mid ev(S \otimes a) = 0, \text{ that is } f(a) = 0, \forall f \in S\}$$

Having associated a topological space to a k -algebra, one might ask when this is a functor. Functoriality has always been a problem in noncommutative geometry. Indeed, a simple B -representation does not have to remain a simple A -representation under restriction of scalars via $\phi : A \longrightarrow B$.

Still, if we define $\text{rep}(A) = A^\circ$, we get functionality for free. If $A \xrightarrow{\phi} B$ is an algebra morphism, we have seen that the dual map maps B° to A° , so we have a morphism

$$B^\circ = \text{rep}(B) \xrightarrow{\phi^*} \text{rep}(A) = A^\circ$$

A coalgebra is the direct limit of its finite dimensional coalgebras, and they correspond under duality to finite dimensional algebras. Hence, ϕ^* is the natural map on finite dimensional representations by restriction of scalars.

The observed failure of functoriality on the level of points translates on the coalgebra-level to the fact that for a coalgebra map $B^\circ \longrightarrow A^\circ$ the coradical $\text{corad}(B^\circ)$ does not have to be mapped to $\text{corad}(A^\circ)$, in general.

However, when $\text{corad}(B^\circ)$ is cocommutative, we *do have* that $\phi^*(\text{corad}(B^\circ)) \subset \text{corad}(A^\circ)$ by [5, Thm. 9.1.4]. In particular, we recover the functor of points in *commutative algebraic geometry*.

Clearly, we still have $\text{corad}(B^\circ) \longrightarrow A^\circ$ in general. This corresponds to the fact that there is always a map $\text{simp}(B) \longrightarrow \text{rep}(A)$.

Next, let us turn to the algebra of functions on $\text{rep}(A)$. By definition we have

$$k[\text{rep}(A)] = A^{\circ*}$$

and we can ask how this algebra relates to the algebra A .

In general, it is *not* true that $A \hookrightarrow A^{\circ*}$. This only holds when A° is dense in A^* in which case the k -algebra is said to be *proper*, see [5, §6.1].

In the commutative case, when A is a finitely generated k -algebra, then A is indeed proper and this is a consequence of the Hilbert Nullstellensatz and the Krull intersection theorem.

When A is noncommutative, this is no longer the case. For example, if $A = A_n(k)$ the *Weyl algebra* over a field of characteristic zero k , then A is simple whence has no twosided ideals of finite codimension. As a result $A^\circ = 0$! As our proposal for the noncommutative affine scheme $\text{rep}(A)$ is based on finite dimensional representations of A , it will not be suitable for k -algebras having few such representations.

3. THE DUAL COALGEBRA A°

In general though, A° is a huge object, so it is very difficult to describe explicitly. In this section, we will begin to tame A° even when A is noncommutative.

In order not to add extra problems, we will assume that $\mathbb{k}$ is separably closed in this section. The general case can be recovered by taking $\text{Gal}(\mathbb{k}/k)$ -invariants (replacing quivers by *species* in the sequel).

Over a separably closed field $\mathbb{k}$ all simple subcoalgebras are full matrix coalgebras $M_n(\mathbb{k})^*$, that is, $M_n(\mathbb{k})^* = \oplus_{i,j} \mathbb{k} e_{ij}$ with $\Delta(e_{ij}) = \sum_{k=1}^n e_{ik} \otimes e_{kj}$ and $\epsilon(e_{ij}) = \delta_{ij}$.

Hence, $\text{corad}(A^\circ) = \oplus_S M_{n_S}(\mathbb{k})^*$ where the sum is taken over all finite dimensional simple A -representations S , each having dimension n_S .

In algebra, one can resize idempotents by Morita-theory and hence replace full matrices by the basefield. In coalgebra-theory there is an analogous duality known as *Takeuchi equivalence*, see [6].

The isotypical decomposition of $\text{corad}(A^\circ)$ as an A° -comodule is of the form $\oplus_S C_S^{\oplus n_S}$, the sum again taken over all simple A -representations. Take the A° -comodule $E = \oplus_S C_S$ and its *coendomorphism coalgebra*

$$A^\dagger = \text{coend}^{A^\circ}(E)$$

then Takeuchi-equivalence (see for example [1, §4, §5] and the references contained in this paper for more details) asserts that A° is Takeuchi-equivalent to the coalgebra $A^\dagger$ which is *pointed*, that is, $\text{corad}(A^\dagger) = \mathbb{k} \text{simp}(A) = \oplus_S \mathbb{k} g_S$ with one *group-like* element g_S for every simple A -representation. Remains to describe the structure of the full basic coalgebra $A^\dagger$.

For a (possibly infinite) quiver $\vec{Q}$ we define the *path coalgebra* $\mathbb{k}\vec{Q}$ to be the vectorspace $\oplus_p \mathbb{k}p$ with basis all oriented paths p in the quiver $\vec{Q}$ (including those of length zero, corresponding to the vertices) and with structural maps induced by

$$\Delta(p) = \sum_{p=p'p''} p' \otimes p'' \quad \text{and} \quad \epsilon(p) = \delta_{0,l(p)}$$

where $p'p''$ denotes the concatenation of the oriented paths p' and p'' and where $l(p)$ denotes the length of the path p . Hence, every vertex v is a group-like element and for an arrow $\textcircled{v} \xrightarrow{a} \textcircled{w}$ we have $\Delta(a) = v \otimes a + a \otimes w$ and $\epsilon(a) = 0$, that is, arrows are skew-primitive elements.

For every natural number i , we define the *extⁱ-quiver* $\overrightarrow{\text{ext}}_A^i$ to have one vertex v_S for every $S \in \text{simp}(A)$ and such that the number of arrows from v_S to v_T is equal to the dimension of the space $\text{Ext}_A^i(S, T)$. With ext_A^i we denote the $\mathbb{k}$ -vectorspace on the arrows of $\overrightarrow{\text{ext}}_A^i$.

The *Yoneda-space* $\text{ext}_A^\bullet = \oplus \text{ext}_A^i$ is endowed with a natural A_∞ -structure [2], defining a linear map (the *homotopy Maurer-Cartan map*, [4])

$$\mu = \oplus_i m_i : \mathbb{k}\overrightarrow{\text{ext}}_A^1 \longrightarrow \text{ext}_A^2$$

from the path coalgebra $\mathbb{k}\overrightarrow{\text{ext}}_A^1$ of the *ext¹-quiver* to the vectorspace ext_A^2 , see [2, §2.2] and [4].

Theorem 1. *The dual coalgebra A° is Takeuchi-equivalent to the pointed coalgebra $A^\dagger$ which is the sum of all subcoalgebras contained in the kernel of the linear map*

$$\mu = \oplus_i m_i : \mathbb{k}\overrightarrow{\text{ext}}_A^1 \longrightarrow \text{ext}_A^2$$

determined by the A_∞ -structure on the Yoneda-space $\text{ext}_A^\bullet$.

We can reduce to finite subquivers as any subcoalgebra is the limit of finite dimensional subcoalgebras and because any finite dimensional A -representation involves only finitely many simples. Hence, the statement is a global version of the result on finite dimensional algebras due to B. Keller [2, §2.2].

Alternatively, we can use the results of E. Segal [4]. Let $S_1, \dots, S_r$ be distinct simple finite dimensional A -representations and consider the semi-simple module $M = S_1 \oplus \dots \oplus S_r$ which determines an algebra epimorphism

$$\pi_M : A \longrightarrow M_{n_1}(\mathbb{k}) \oplus \dots \oplus M_{n_r}(\mathbb{k}) = B$$

If $\mathfrak{m} = \text{Ker}(\pi_M)$, then the $\mathfrak{m}$ -adic completion $\hat{A}_{\mathfrak{m}} = \varprojlim A/\mathfrak{m}^n$ is an augmented B -algebra and we are done if we can describe its finite dimensional (nilpotent) representations. Again, consider the A_{∞} -structure on the Yoneda-algebra $\text{Ext}_A^{\bullet}(M, M)$ and the quiver on r -vertices $\overrightarrow{\text{ext}}_A^1(M, M)$ and the homotopy Maurer-Cartan map

$$\mu_M = \oplus_i m_i : \overrightarrow{\text{ext}}_A^1(M, M) \longrightarrow \text{Ext}_A^2(M, M)$$

Dualizing we get a subspace $\text{Im}(\mu_M^*)$ in the path-algebra $\overrightarrow{\text{ext}}_A^1(M, M)^*$ of the dual quiver. Ed Segal's main result [4, Thm 2.12] now asserts that $\hat{A}_{\mathfrak{m}}$ is Morita-equivalent to

$$\hat{A}_{\mathfrak{m}} \underset{M}{\sim} \frac{(\overrightarrow{\text{ext}}_A^1(M, M)^*)^{\wedge}}{(\text{Im}(\mu_M^*))}$$

where $(\overrightarrow{\text{ext}}_A^1(M, M)^*)^{\wedge}$ is the completion of the path-algebra at the ideals generated by the paths of positive length. The statement above is the dual coalgebra version of this.

REFERENCES

- [1] William Chin, *A brief introduction to coalgebra representation theory*, in "Hopf Algebras" M. Dekker Lect. Notes in Pure and Appl. Math. (2004) 109-132. Online at <http://condor.depaul.edu/~wchin/crt.pdf>
- [2] Bernhard Keller, *A-infinity algebras in representation theory*, Contribution to the Proceedings of ICRA IX, Beijing (2000). Online at <http://www.math.jussieu.fr/~keller/publ/art.dvi>
- [3] Maxim Kontsevich and Yan Soibelman, *Notes on A_{∞} -algebras, A_{∞} -categories and non-commutative geometry I*, arXiv:math.RA/0606241 (2006)
- [4] Ed Segal, *The A_{∞} deformation theory of a point and the derived category of local Calabi-Yaus*, math.AG/0702539 (2007)
- [5] Moss E. Sweedler, *Hopf Algebras*, monograph, W.A. Benjamin (New York) (1969)
- [6] M. Takeuchi, *Morita theorems for categories of comodules*, J. Fac. Sci. Univ. Tokyo 24 (1977) 629-644

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ANTWERP, MIDDELHEIMLAAN 1, B-2020 ANTWERP (BELGIUM), lieven.lebrun@ua.ac.be